

Inceptova[®] FSH

HIGHLY PURIFIED

Urofollitropin for injection



Description^{1,2}

Inceptova FSH is a product containing a highly purified preparation of human follicle stimulating hormone (hFSH) extracted from the urine of postmenopausal women.

Composition

Each vial of sterile freeze-dried product contains:

- Urofollitropin B.P. equivalent to the activity of
 - FSH BP 75 IU
 - Mannitol IP q.s.
 - Potassium Dihydrogen Phosphate BP q.s.
 - Dipotassium Hydrogen Phosphate BP q.s.
- Urofollitropin B.P. equivalent to the activity of
 - FSH BP 150 IU
 - Mannitol IP q.s.
 - Potassium Dihydrogen Phosphate BP q.s.
 - Dipotassium Hydrogen Phosphate BP q.s.

Indications²

Inceptova FSH provides ovarian follicular growth and maturation in females who do not have primary ovarian failure. Also stimulates spermatogenesis in males (off-label use).

Dosage¹

Adults – at first, 225 IU injected under the skin once a day. Doctor will adjust the dose after 5 days, if needed. However, the dose is usually not more than 450 IU per day. Do not use FSH and LH for more than 20 days.

Special Precautions and Warnings²

- Do not use **Inceptova FSH**, if you have primary ovarian failure, abnormal vaginal bleeding, uncontrolled thyroid or adrenal gland disorders, an ovarian cyst, breast cancer, uterine or ovarian cancer, a tumor of pituitary gland or hypothalamus or infertility that is not caused by lack of ovulation.
- Do not use **Inceptova FSH**, if pregnant.

Drug Interactions²

There are no known significant interactions.

Undesirable Effects²

- **Less Common:** Back pain, breast tenderness, feeling of warmth, redness of the face, neck, arms, and occasionally, upper chest, menstrual changes, muscle aches and pains and unusual tiredness or weakness.
- **Rare:** Abdominal or stomach pain (severe), bloating (moderate to severe), chest pain or trouble breathing, decreased amount of urine, feeling of indigestion, general feeling of discomfort or illness, headache, severe and throbbing, nausea, vomiting or diarrhea (continuing or severe), pain or swelling in the arms or legs, pelvic pain (severe), severe cramping of the uterus, shortness of breath or wheezing, swelling of the lower legs and weight gain (rapid).

Contraindications²

Use during pregnancy is contraindicated.

Hypersensitivity; primary ovarian failure as indicated by a high FSH level; uncontrolled nongonadal endocrinopathies (e.g. thyroid, adrenal, pituitary); pituitary or hypothalamic tumors; abnormal uterine bleeding of undetermined origin; ovarian cyst or enlargement not due to PCOS; pregnancy.

Inceptova FSH is also contraindicated with sex hormone-dependent tumors of the reproductive tract and accessory organs.

Special Precautions for Storage²

Store at room temperature away from moisture, heat and light.

Do not freeze.

References

1. <https://www.drugs.com/cons/follicle-stimulating-hormone-and-luteinizing-hormone-intramuscular-subcutaneous.html>
2. <https://www.drugs.com/ppa/menotropins.html>

™ Trade Mark

Marketed by:



Human Biologicals Institute
(A Division of Indian Immunologicals Limited)

Manufactured by:

Sanzyme (P) Ltd.
Plot No. 8, Sy. No. 542, Koltur (V),
Shamirpet (M), Medchal-Malkajgiri (Dist.),
500101, Telangana, INDIA.